

## STUDY ON THE ATTACK OF *PSEUDOCERCOSPORELLA* *HERPOTRICHOIDES* ON WHEAT STEM BASE UNDER CONDITIONS FROM TIMISOARA DIDACTIC RESEARCH STATION

Diana VELIMIROVICI, Adrian BORCEAN

*Banat's University of Agricultural Sciences and Veterinary Medicine, Faculty of Agricultural Sciences, Timisoara, Aradului Street, no. 119, RO-300645, Romania,*  
Corresponding author: vlmiana@yahoo.com

**Abstract:** Research aim was to test the tolerance of a range of wheat cultivars to infectious pressure of the pathogen *Pseudocercospora herpotrichoides* in climatic conditions from Timisoara Didactic Research Station. Stage of research is being characteristic for partial interpretation of experimental data after the first experimental year, in preparation the doctoral thesis. Materials and methods. Experimental field was located in climatic conditions from Timișoara. Technology was the standard applied to this area. The experience was bifactorial. First factor was the cultivar with three graduations (Ciprian, Arieșan and Kristina). Factor B was the nitrogen amount with three graduations ( $N_{50}$ ,  $N_{100}$  and  $N_{150}$ ) all three doses were applied on a constant background elements of  $P_{60}$   $K_{60}$ . The novelty is relatively high, work providing important data for agricultural practice in the experimental area, taking in consideration the implications of fungus *Pseudocercospora herpotrichoides* in yield amount and harvesting problems in some years. Achievements stage in this field. In this work were carried out research including one of authors, known as the reference and possible reactions of the pathogen depending on local biocoenosis factors, including the competition between stem base pathogens. Limits of the research are that data from the fungus *Pseudocercospora herpotrichoides* are just after one year bonitation. Practical implications of the research consisted of playing a part of a complex study of strategy in the of wheat protection reference experience. The originality of the work comes from the fact that data are relevant in view of cultivars behaviour under specific conditions. These data are only part of the data submitted for interpretation in the preparation of the doctoral experience of the author. Importance of the paper became from bringing in front of the specialists of one of the topics in the experiences of the author's doctoral preparing, to evaluate the accuracy of techniques addressed. The relevant data, experimental results give an overview of pathogen behavior in relation to experimental conditions.

**Key words:** *Pseudocercospora herpotrichoides*, eye spot of wheat stem, fertilizers

### INTRODUCTION

Between March and end of May 2009 we have made a number of observations in the wheat experimental of field crops discipline from Didactic Research Station. It was very interesting to see the differences between hybrids to the attack of fungus *Pseudocercospora herpotrichoides* and, as a part of a larger study concern the main pathogens of the stem base and roots of wheat behavior in the climate conditions from Timisoara.

### MATERIAL AND METHOD

Experimental conditions where those from Didactic Research Station from Timișoara and consist from a chernozem soil and ecological conditions. In the figures below are described temperature (figure 1) and rain (figure 2) from the year 2009 and the multiannual average for this two whether parameters available for the experimental field.

Assessing pathogen population dynamics was done by statistic analysis of attack degree, calculated as synthetic indicator on the base of the readings performed on the experimental field of frequency and intensity of attack in the interval between March and May

2009, when the pathogen of the root and stem base of the wheat were monitored at the Didactic Station of the University of Agricultural Sciences and Veterinary Medicine of the Banat in Timișoara in a bi-factorial experiment.




Figure 1. Monthly temperature averages, compared with long term averages recorded at Timișoara Meteorological Station




Figure 2. Monthly rainfall water amount, compared with long term averages recorded at Timișoara Meteorological Station

The factors we monitored during the three experimental years were the cultivated cultivar and amount of nitrogen applied. The two factors we monitored strongly influence the appearance and development of attack by pathogens, as follows: cultivar by resistance or degree of tolerance of cultivars to attack and nitrogen amount by the physiological influence on the length of different vegetation period and on the influence of physiological vigour of the plants and the easiness the pathogen is transmitted.

The first factor, experimental cultivar, had three graduations: Ciprian, Arieşan and Kristina. The second factor, the nitrogen amount also with three graduations, on the background of constant phosphorus and potassium amount: N<sub>100</sub> P<sub>60</sub> K<sub>60</sub>, N<sub>150</sub> P<sub>60</sub> K<sub>60</sub> and N<sub>200</sub> P<sub>80</sub> K<sub>80</sub>.

Statistic calculation was done after the model of experiment with two factors.

Attack frequency and intensity were read from the experimental trials in two different vegetation stages:

- first reading is on the stage when the second node of the stem is at least 2 cm above first node, the stage 32 on BBCH scale
- second reading when plants are on early milk stage, the stage 73 on BBCH scale.

### RESULTS AND DISCUSSIONS

All the observations figures concerning attack frequency and intensity from the field are in table 1. The attack degree calculated based on frequency and intensity of attack are also in table 1. Considering the attack degree is a synthetic indicator of attack frequency and intensity of attack and this is the reason why we will refer at it for statistic calculation.

In the year 2009, frequency registered normal values, but the intensity of attack have very low values than frequency and as it can be observed, the higher values was in the trials where was applied the largest amount of nitrogen.

Table 1.

Field readings for frequency and intensity and calculated attack degree of fungus *Pseudocercospora herpotrichoides* for the experimental variants in the year 2009.

Nr. crt.	Hibridul	Agrofondul	Frecvența (%)				Intensitatea (%)				Gradul de atac			
			R1	R2	R3	x	R1	R2	R3	x	R1	R2	R3	x
1	Ciprian	N <sub>100</sub> P <sub>60</sub> K <sub>60</sub>	10	5	10	8,3	1	5	1	2,3	0,1	0,25	0,1	0,15
		N <sub>150</sub> P <sub>60</sub> K <sub>60</sub>	10	12	17	13	5	3	5	4,3	0,5	0,36	0,85	0,57
		N <sub>200</sub> P <sub>60</sub> K <sub>60</sub>	15	20	14	16,3	10	5	5	6,6	1,5	1	0,7	1,06
2	Arieşan	N <sub>100</sub> P <sub>60</sub> K <sub>60</sub>	10	15	14	13	1	5	5	3,6	0,1	0,75	0,7	0,51
		N <sub>150</sub> P <sub>60</sub> K <sub>60</sub>	17	23	15	18,3	10	5	10	8,3	1,7	1,15	1,5	1,45
		N <sub>200</sub> P <sub>60</sub> K <sub>60</sub>	10	26	18	18	10	20	25	18,3	1	5,2	4,5	3,56
3	Kristina	N <sub>100</sub> P <sub>60</sub> K <sub>60</sub>	15	10	12	12,3	5	5	5	5	0,75	0,5	0,6	0,61
		N <sub>150</sub> P <sub>60</sub> K <sub>60</sub>	15	15	22	17,3	5	10	5	6,6	0,75	1,5	1,1	1,12
		N <sub>200</sub> P <sub>60</sub> K <sub>60</sub>	23	16	21	20	15	10	15	13,3	3,45	1,6	3,15	2,73
4	Average	N <sub>100</sub> P <sub>60</sub> K <sub>60</sub>	12	12,4	13,4	12,6	4,4	4,6	4,2	4,4	0,59	0,57	0,59	0,58
		N <sub>150</sub> P <sub>60</sub> K <sub>60</sub>	13,4	17	16,4	15,6	6,2	5,6	6	5,9	0,91	0,95	0,97	0,94
		N <sub>200</sub> P <sub>60</sub> K <sub>60</sub>	15	20	16,4	17,1	9,2	9	12	10	1,55	1,94	2,11	1,86

Infectious pressure of fungus *Pseudocercospora herpotrichoides* is shown by the fungus by this values of frequency and intensity because all the readings was done after infections in natural conditions and after a few years with low rate of rains. The quantity of water from rain is lower then multiannual average (figure 2) and the rain distribution is deficient because the water was collected in just a few rains.

The biological material used in the experience, prove to have a nice diversity, based on the variation of the frequency and intensity values. In the same time, if the variation of frequency and intensity show one more time that the tolerance of varieties is an individual propriety, it have a strict dependency on the weather conditions and technology applied.

Statistic calculation performed for attack degree data(table 2), point out that the differences of varieties average are under the limit of significance and it can be explained, if we take a closer look at data from table 1. and also to the figure 2. There is shown that because of the lack of the rains in the first five month of the year 2009, the experimental condition wasn't so favorable to diseases. Under those conditions all varieties registered at least a normal frequency of attack and lower intensity of attack of fungus *Pseudocercospora herpotrichoides*. The result was a very balanced situation on the attack degree, proved by the luck of differences on the attack degree averages regarding to the experimental average and this was the defining behavior of tested varieties on fungus *Pseudocercospora herpotrichoides* attack.

Table 2  
Statistic interpretation data for attack degree of fungus *Pseudocercospora herpotrichoides*.  
in the year 2009

Factor A Variety	Factor B - Fertilizers			Averages of Factor A	Differences	Significance
	N <sub>100</sub> P <sub>80</sub> K <sub>80</sub>	N <sub>150</sub> P <sub>80</sub> K <sub>80</sub>	N <sub>200</sub> P <sub>80</sub> K <sub>80</sub>			
Ciprian	0,15	0,57	1,06	0,59	- 0,68	-
Arieşan	0,51	1,45	3,56	1,84	0,57	-
Kristina	0,61	1,12	2,73	1,49	0,22	-
Average	0,58	0,94	1,86	1,27	Witness	-

DL 5% = 0,89      DL 1% = 1,34      DL 0,1% = 2,16

Factor B - Fertilizers	N <sub>100</sub> P <sub>80</sub> K <sub>80</sub>	N <sub>150</sub> P <sub>80</sub> K <sub>80</sub>	N <sub>200</sub> P <sub>80</sub> K <sub>80</sub>
Averages of Factor B	0,58	0,94	1,86
Differences	Witness	0,35	1,28
Significance	-	-	***

DL 5% = 0,65      DL 1% = 0,90      DL 0,1% = 1,24

The nitrogen prove to have a stimulating effect on eyespot , the experimental averages has no significance for 150 kg/ha. Growing the nitrogen dosage to 200 kg /ha lead to an increase of fungus attack which registered a very significant difference (table 2).

### CONCLUSIONS

- Frequency and intensity of fungus *Pseudocercospora herpotrichoides* attack have a very low variation on all varieties which prove to have a very slight tolerance to this fungus, specially when there are technology factors that stimulate the fungus attack.
- Nitrogen application increases the disposition of the plants to be infected by eyespot by extension of the period of maximum sensitivity.
- Attack degree registered a very low difference between varieties, the values was under the significance threshold reported to experimental average.
- Regarding to the effect of nitrogen, it is very clear that the increasing of nitrogen dosage conduct to an increase of attack degree wich registry a distinct significant value of difference on the nitrogen amount of 150 kg/ha and a very significant difference on the nitrogen amount of 200 kg/ ha.

### BIBLIOGRAPHY

1. BĂLTEANU GH., BÎRNAURE V., SALONTAI AL., BORCEAN I., VASILICĂ C., Fitotehnie, Ed. Didactică și Pedagogică București, 515 pag., 1991
2. BORCEAN I., TABĂRĂ V., PÎRȘAN P., DEMENESCU T., BORCEAN A., Cercetări privind stabilirea principalelor verigi tehnologice la porumb în condițiile Banatului, *Lucrări Științifice, U.A.Iași, Vol. XXXIV*, pag 157-163, 1992.
3. BORCEAN I., TABĂRĂ V., PÎRȘAN P., DAVID GH., POP GEORGETA, BORCEAN A., ȘUVEȚI SIMONA, CIULOI ANCA., BORUGĂ L. Comportarea a 40 hibrizi de porumb în condițiile cernoziomului de la Timișoara, *Simpozin-Cultura plantelor de câmp U.S.A.M.V.B. Timișoara*, pag 3-9, 1996.
4. DAVID GH., BORCEAN A., Sweet corn – A crop plant with high perspectives on the Western Plain of Romania, University of Agricultural Sciences College of Agriculture Mezötúr, 1999.
5. BORCEAN I., MIRCOV V., IMBREA F., BORCEAN A., Preliminary results concerning the influence of electromagnetic treatments at maize, *Universitatea Novi Sad, Biofizika u poljoprivrednoj proizvodnji*, pag. 187-190, Novi Sad 2002
6. BORCEAN I., DAVID GH., BORCEAN A., IMBREA F., BOTOȘ L., On the behaviour of some new maize hybrids in the conditions of brown luvisc soils in the hill area of the Banat, *Lucrări Științifice Facultatea de Agricultură, U.S.A.M.V.B. Timișoara, Vol XXXIV*, pag 187-192, 2002.
7. BORCEAN A., DAVID GH., MOLNAR L., Observații privind comportarea unui sortiment de hibrizi de porumb la atacul ciupercii *Ustilago maydis* în condițiile anului 2006 de la SD a USAMVB Timișoara. *Trends in European agriculture development, Faculty of Agriculture Timisoara and Faculty of Agriculture Novi Sad International Symposium, 2007*
8. BORCEAN I., BORCEAN A., 2004, *Cultura și protecția integrată a cerealelor, leguminoaselor și plantelor tehnice*, Editura de Vest, 509 pag.
9. BORCEAN I., ȚĂRĂU D., BORCEAN A., DAVID GH., BORCEAN EUGENIA, *Fitotehnia și protecția culturilor de câmp*, Editura de Vest, 198 pag.
10. BORCEAN I., DAVID GH., BORCEAN A., *Tehnici de cultură și protecție a cerealelor și leguminoaselor*, Editura de Vest, 346 pag.
11. POPESCU GH. – *Fitopatologie*, Ed. Mirton, Timișoara, 1998
12. M. HATMANU, I. BOBEȘ, AL. LAZĂR, C. GHEORGHIȘ, C. GLODEANU, V. SEVERIN, Corina Tușa – *Fitopatologie*, Ed. Didactică și Pedagogică, București, 1989, 468 pag.
13. NIȚĂ SIMONA, BORCEAN A., Study regarding some varieties of Zea mays everta Stuart which grow under the pedo-climatic conditions in Almaj Depression, „60 de ani de învățământ superior în Oltenia”, *Annales of University of Craiova, Vol XXXVII / A 2007*, pag. 664-666, 2007
14. PÎRȘAN P., DAVID GH., BORCEAN A., WAUTERS S., Cercetări cu privire la studiul unui sortiment de hibrizi proveniți din vestul Europei în condițiile de la Timișoara, *Lucrări Științifice, U.S.A.M.V.B. Timișoara, Vol XXX*, 1998.