

**PRELIMINARY STUDIES ON THE VOLATILE OIL CONTENT OF
HERBAL AND THE MENTHOL PEPPERMINT (*MENTHA PIPERITA L.*)
UNDER THE INFLUENCE OF FERTILIZATION IN MINERAL LOVRIN**

O. PĂTRUI, V. TABĂRĂ

*Banat's University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine,
119, Calea Aradului, 300645 Timișoara, Romania
E-mail: patruiovidiu@yahoo.com*

Abstract: *Mint is one of the oldest herbs known. Mint is an herb well known for its medicinal properties, but also for refreshing flavor. Given the fact that fertilization is one of the measures which, properly applied, lead to increased production of foil, production of volatile oil and the oil content and hence its quality, more research is carried out lately have focused on finding optimal formulas for this. The volatile oil is the most valuable component of the composition of mint leaves (foil). The quality of healing depends on the quality of the finished product that is found in almost all health and beauty products. Calculation of volatile oil production has been achieved in relation to film production in mint. The essential oil obtained from the leaves of *Mentha piperita L.*, has numerous applications in medicine due to antibacterial properties, antiviral and cytotoxic essential oil, fragrance oil and antioxidants properties make it to be used in food and beverage industry. In England, as in the rest of Europe, peppermint is used almost exclusively for sweet and sweet liqueurs where her pungent fresh and refreshing counterbalance to the sweetness of sugar properties. For these purposes, the use of pure essential oil is preferred to avoid the bitter astringent notes of peppermint leaves. Peppermint freshness fits extremely well with chocolate. Peppermint ice cream is delicious on a warm summer day, because it uses soft properties of menthol. Mint is grown in many countries in*

*Europe and Central and West Asia for the production of menthol, which is necessary in pharmaceutical preparations. In most of these countries entered peppermint local cuisine, replacing the local varieties mint. The study aims to enrich knowledge on the role of characteristics of cultivation technologies mint (*Mentha piperita L.*), to achieve improved their performance, to achieve a high production of essential oil of a higher quality, thus the study of the possibility expansion of cultivation areas are mint. Experience has been located on land belonging in Lovrin. Experience is the type bifactorial with annual repetition. The biological material used was kind Column, as certified seed. Organic and inorganic fertilizers applied peppermint culture, increased production of essential oil and menthol in the foil. The content of active substances, the determination of volatile oil content of mint and menthol, was monitored in the laboratories S.C. Fares Vital Bio Laboratories LLC Drastic, (ISO 9001: 2000). The results of this study are part of a PhD program, having as theme: "Development of cultivation technology of mint (*Mentha piperita L.*) in conditions of Lovrin". The results of this study are part of a PhD program, having as theme: "Development of technology for cultivation of mint (*Mentha piperita L.*) in conditions of Lovrin" under the leadership of the distinguished university professor Valeriu Tabără.*

Key words: *peppermint, volatile oil, herba, menthol.*

INTRODUCTION

Peppermint (*Mentha piperita L.*) is one of the most popular and widely consumed aromatic and medicinal plants by people. The reason that peppermint is so popular is that it spreads a great smell nice, but mostly because it is an excellent herb tea that gives a great taste. Being a plant originating in northern Europe, is found in spontaneous, instead it is cultivated on large areas in many parts of the world. Peppermint oil contains high levels of essential active menthol.

The volatile oil of peppermint and the other components of the plant have stomahică

action, carminative, choleric, cardiotoxic, analgesic.

Menthol is widely used in the preparation of toothpaste and mouth washes because of soft properties, antiseptics and the correction of taste and smell.

Peppermint essential oil is widely used in food, the preparation of liqueurs, candies, which prints an odor and a pleasant.

Proper fertilization with manure, which is the best fertilizer for mint (especially on poor soils), depending on soil applied doses of 20-40 t/ha (incorporated by plowing), it is very important not only for quality production, but also to maintain and improve the productive potential of soil. Herba and foil production is influenced by the technology applied (especially fertilization) that determines the quality and increase production.

MATERIAL AND METHODS

Experience has been placed in the experimental field of research Lovrin resort. The experimental field was located on a chernozem soil type. The experiments were bifactorial type, to repeat the concluding annual cycle so that the experimental field in the first year we have experience, second year and third year. Variety Columna used in experiments.

FACTOR A - organic fertilization - 20 t / ha

- a1: N₀P₀K₀
- a2: N₆₀P₆₀K₆₀
- a3: N₉₀P₉₀K₉₀
- a4: N₁₂₀P₉₀K₉₀
- a5: N₁₂₀P₉₀K₉₀ + foliar fertilization

FACTOR B – unfertilized organic

- a1: N₀P₀K₀
- a2: N₆₀P₆₀K₆₀
- a3: N₉₀P₉₀K₉₀
- a4: N₁₂₀P₉₀K₉₀
- a5: N₁₂₀P₉₀K₉₀ + foliar fertilization

RESULTS AND DISCUSSIONS

Volatile oil and menthol content of the herbal to mint (*Mentha piperita L.*) in the experimental field of Lovrin.

Results of the content of volatile oil obtained from herba production in mint (*Mentha piperita L.*) under the influence of factor A - the mineral and organic fertilized Lovrin are shown in Figure 1.




Figure 1. Changes in the mean content of volatile oil obtained from the mint herba under the influence of organic fertilization (factor A), Lovrin

It was found that the volatile oil is formed in the plant, depending on the origin of some genetic conditions, time of harvest, the leaves on the stem position. The analysis results shows that there is a high variability of essential oil content of mint foil at the 5 variants studied. This variability is within the limits, 1.5% oil only organic and mineral fertilized variant 1.74% at 120 N. version.

Of the five variants studied 3 to 1 oil content of over 1.6% (, N90P90K90, N120P90K90 and N120P90K90 + FF).

Of the variants studied, four exceed the oil content of only organic mineral fertilized variant of 1.5%.

Variation of volatile oil content and its components that differ from herba of mint, depending on the time of harvesting their geographical location.

Table 1

Influence of organic and mineral fertilization on the production of essential oil from the herba at the mint, the Lovrin

Nr.crt.	A – organic/ mineral fertilizer	Crops kg/ha	%	Diference kg/ ha	Significance
1.	a1 N0P0K0	180	100	-	
2.	a2 N60P60K60	218	121	38	
3.	a3 N90P90K90	269	149	89	xxx
4.	a4 N120P90K90	298	166	118	xxx
5.	a5 N120P90K90 + FF	304	169	124	xxx

DL5% = 43 kg/ha; DL 1% = 61 kg/ha; DL 0,1% = 88 kg/ha.

Production for the factor A given product is a result harvest - oil content.

The higher oil production from Lovrin under the influence of mineral and organic fertilization is achieved in variants N120P90K90 + FF - 304kg/ha oil N120P90K90 - 298kg/ha oil N90P90K90 - 269 kg / ha oil, and N60P60K60 - 218 kg / ha oil. It appears that the 5 variants studied, four oil production achieved over 250 kg / ha and oil that are statistically very significant. The smallest production of essential oil of peppermint is recorded on film from mineral fertilized variant only organic NOP0K0 with a value of 180 kg / ha oil.

The content of volatile oil obtained from herba yield in peppermint (*Mentha piperita L.*) under the influence of **factor B** to Lovrin.

Analysis of volatile oil content of the herba at the mint produced by sowing mineral fertilization variants under the influence of climatic conditions Lovrin reveals that four of the five variants have a higher oil content in the foil version NOP0K0 unfertilized (1,36%) .




Figure 2. Influence of fertilization on oil content of herba yield in peppermint (*Mentha piperita L.*) in Lovrin

The high content of essential oil of herba mint under the influence of sowing mineral fertilization performed on variants N60P60K60 - 1.49%; N90P90K90 - 1.56%; N120P90K90 - 1.66% and N120P90K90 + FF - 1.56 %. The lowest oil content was recorded in the variant fertilized N0P0K0 - 1.36%.

If we compare the content of volatile oil of herba in mint planted under the influence of mineral and organic fertilization and sowing mineral fertilization influence under the influence note that by sowing mineral fertilization, the contents of volatile oil of herba, all variants will be reduced.

Production of essential oil of herba Mentor under the influence of mineral fertilization is presented in Table 2.

Table 2

Influence of mineral fertilization on the yield of essential oil of peppermint in the herba at the experimental field Lovrin

Nr.crt.	B mineral fertilizer	Crops kg/ha	%	Diference kg/ ha	Significance
1.	b1 N0P0K0	181	100	-	
2.	b2 N60P60K60	221	122	40	xxx
3.	b3 N90P90K90	283	156	102	xxx
4.	b4 N120P90K90	291	161	110	xxx
5.	b5 N120P90K90 + FF	280	155	99	xxx

DL5%= 19 kg/ha; DL 1% = 28 kg/ha; DL 0,1% = 40 kg/ha.

Analyzing the results of the oil production under the influence of 5 different sowing mineral fertilization of Lovrin conditions shows the following:

- production of essential oil of mint is dependent herba in the production of herba,
- production of herba determine the level and volatility of oil production,
- volatile oil content falls under the influence of sowing mineral fertilization,
- the difference between variants is higher in the production of essential oil of the peppermint herba sown under the influence of organic fertilization.

Such variants N60P60K60 and N90P90K90, N120P90K90, N120P90K90 + FF increases the production of essential oil from the film are that 40,102,110 99 kg / ha and oil are statistically very significant. Highest yields of volatile oil of herba at Spearmint is recorded on the N90 variants with mineral fertilizer - 283 kg / ha oil and N120 - 291 kg / ha oil.

Menthol content in the foil at the mint under the influence of mineral and organic fertilization.

Chief component of the essential oil of peppermint is menthol foil. After the parts were removed from menthol oil, oil is marked as (dementolizat corrected).




Figure 3. Menthol content of mint herba at the Lovrin under the influence of organic and mineral fertilization

Menthol is the most valuable component of the composition of the herba of mint.

The analysis results shows that there is a high variability of herba menthol content in peppermint. This variability is within the limits of 28.27% menthol in organic fertilized variant only - N0P0K0 and 30.59% for the variant N120P90K90 + FF.

Of the five variants studied only tdoau have a menthol content of over 30% (FF + N120P90K90 - and N120P90K90 30.59% - 30.11%).

The lowest amount of menthol content of the film under the influence of organic and mineral fertilization conditions in Lovrin N0P0K0 are recorded version - 28.27%.

Menthol content in the herbal at the mint under the influence of mineral fertilization. Menthol content in peppermint yield of the herba under influența mineral fertilization is presented in Figure 4.




Figure 4. Influence of mineral fertilization on the content of menthol in mint foil at the conditions of Lovrin

Analysis of the menthol content in peppermint herba obtained under the influence of mineral fertilization conditions in Lovrin point out that four of the variants studied have a higher menthol content unfertilized variant (23.94%).

The high content of menthol in mint foil at the conditions of Lovrin performed on variants N120P90K90 - 26.08% and N120P90K90 + FF - 25.88%.

Under the menthol content of the film are variants with variants 120N 90N - 25.8% menthol, 60N - 24.14% and unfertilised variant - 23.94% menthol.

A single line has a menthol content of greater than 26% (N120P90K90 - 26.08%). It is noted that the five variants studied safflower new study, only 3 to 1 over 25% menthol content.

If we compare the content of menthol in the herba mint obtained under the influence of mineral and organic fertilization and menthol content from the herba obtained under the influence of mineral fertilization, we find that under the influence of mineral fertilization, the content of menthol mint in the foil is reduced by 4%.

CONCLUSIONS

In the Lovrin experimental field was recorded good results in oil and menthol content of mint herba at harvest.

Climatic conditions during sowing and harvesting have largely positive influence on the level of the foil at the mint crop.

Volatile oil content and oil production in mint herba are different depending on the influence of organic and mineral fertilization.

In the variant sown under the influence of organic and mineral fertilization were obtained the highest values of volatile oil content, menthol and oil production.

It is noteworthy that mineral fertilization to have a production decreases in the content of volatile oil, menthol and peppermint oil production of each variant, compared with the content of volatile oil, menthol and mint oil production made under the influence of organic fertilizers and minerals.

If we compare the content of menthol in the herba mint obtained under the influence of mineral and organic fertilization and menthol content from the herba obtained under the influence of mineral fertilization, we find that under the influence of mineral fertilization, the content of menthol mint in the foil is reduced by 4%.

BIBLIOGRAPHY

1. BĂLTEANU GH., 1993 - Plante medicinale în fitotehnie, Ed. Ceres București
2. CIULEI I., GRIGORESCU EM., URSULA STĂNCULESCU, 1993– Plante medicinale, fitochimie și fitoterapie, Tratat de farmacognozie, Vol. II, Ed. Medicală, București
3. COCIU E., RACZ G., 1962 - Plante medicinale și aromatice, Ed. Acad. Române București
4. CROTEAU R, MARTINKUS C (1979) Metabolism of monoterpenes: demonstration of (1) - neomenthyl-b-d-glucoside as a major metabolite of (2) - menthone in peppermint (*Mentha piperita*). Plant Physiol 64: 169–175
5. MUNTEAN L.S. et.al., 2007 – Tratat de plante medicinale cultivate și spontane, Ed. Risoprint, Cluj Napoca
6. SAULESCU N. A., SAULESCU N., 1967 - Câmpul de experiență. Ed. Agrosilvică, București